



E-ISSN: 2706-9575  
P-ISSN: 2706-9567  
IJARM 2020; 2(1): 41-45  
Received: 26-01-2020  
Accepted: 28-02-2020

**Dr. Sanjay Kumar**  
Assistant Professor,  
Department of Medicine, Shri  
Ram Murti Smarak Institute  
of Medical Sciences, Bareilly -  
Nainital Rd, Bhoji Pura, Uttar  
Pradesh, India

**Dr. Anita Kumari**  
Professor, Department of  
Paediatrics, Shri Ram Murti  
Smarak Institute of Medical  
Sciences, Bareilly - Nainital  
Rd, Bhoji Pura, Uttar  
Pradesh, India

**Corresponding Author:**  
**Dr. Anita Kumari**  
Professor, Department of  
Paediatrics, Shri Ram Murti  
Smarak Institute of Medical  
Sciences, Bareilly - Nainital  
Rd, Bhoji Pura, Uttar  
Pradesh, India

## Spectrum of bone disorders in chronic kidney disease patients: A teaching hospital based study in North India

**Dr. Sanjay Kumar and Dr. Anita Kumari**

**DOI:** <https://doi.org/10.22271/27069567.2020.v2.i2a.42>

### Abstract

**Background:** Chronic kidney disease (CKD) is a progressive loss in renal function which involves in deterioration in mineral homeostasis with disruption of normal serum and tissue concentration of phosphorus and calcium. Also changes in circulating levels of hormones parathyroid hormone (PTH), calcitriol (1, 25(OH)<sub>2</sub> D), and fibroblast growth factor-23 (FGF-23). Here our aim is to study the prevalence of markers associated with MBD in CKD stage 3-5 patients. Patients with CKD stage 3-5 were included in this observational study with all necessary parameter. X-RAY abdomen and echocardiography was done to look for evidence of vascular and valvular calcification respectively. Statistical analysis was done using SPSS software. A total of 175 patients (132 males, 43 females)(M:F = 3:1) were included in this study with a mean age of 50.54 years. Among CKD stages 3 to 5, the prevalence of hypocalcemia was 21.5%, 33.9% & 48.9%, hyperphosphatemia was 11.1%, 25.5% & 63%, hyperparathyroidism was 48.1%, 67.3% & 89.1%, high total alkaline phosphatase was 0%, 5.9% & 45.7%, low 25-OH-vit D was 59.2%, 70.6% & 79.4% respectively. Low 25 (OH) D levels, hyperparathyroidism, and hyperphosphatemia were the noticeable markers of CKD-MBD in our patients. Mineral bone disorder are common in CKD patients which start in early CKD stages & worsen with disease progression that causes morbidity and decreased quality of life.

**Keywords:** CKD, MBD, hyperparathyroidism, hyperphosphatemia, hypocalcemia

### Introduction

Chronic kidney disease (CKD) has a prevalence of 5-10% of the world population <sup>[1]</sup>. Patients with CKD have high rates of total and cardiovascular morbidity and mortality.

Abnormalities in levels of mineral metabolites and regulators, such as calcium, phosphate, parathyroid hormone, fibroblast growth factor-23 and vitamin D have been linked to the progression of cardiovascular disease and poor outcomes <sup>[2]</sup>. These mineral and endocrine capacities are basically significant in the guideline of both starting bone formation during development and bone structure & function all through adulthood (modeling and remodeling of bone) <sup>[3]</sup>.

As kidney function declines, there is a progressive deterioration in mineral homeostasis with a disruption of normal serum and tissue concentrations of phosphorus and calcium and changes in circulating levels of hormones. These include parathyroid hormone (PTH), 25-hydroxy vitamin D, 1, 25-dihydroxyvitamin D and other vitamin D metabolites, fibroblast growth factor-23 (FGF-23), and growth hormone.

The ability of the kidneys to appropriately excrete a phosphate load is diminished when GFR falls below 60, leading to hyperphosphatemia, elevated PTH, decreased 1,25(OH)<sub>2</sub>D with associated elevations in the levels of FGF- 23. The conversion of 25(OH) D to 1, 25(OH) 2 D is impaired, reducing intestinal calcium absorption and increasing PTH. The kidneys fail to respond adequately to

PTH, which normally promotes phosphaturia and calcium absorption, or to FGF-23, which also enhances phosphate excretion. In addition, there is evidence at the tissue level of a down regulation of vitamin D receptor and of resistance to the actions of PTH. Therapy is generally focused on correcting biochemical and hormonal abnormalities in an effort to limit their consequences. The mineral and endocrine functions disrupted in CKD are critically important in the regulation of both initial bone formations during growth (bone modeling). As a result, bone abnormalities are found almost universally in patients with CKD requiring

dialysis (stage 5D), and in the majority of patients with CKD stages 3-5. More recently, there has been an increase concern of extra skeletal calcification that may result from the deranged mineral and bone metabolism of CKD and from the therapies used to correct these abnormalities [4].

Numerous cohort studies have shown associations between disorders of mineral metabolism and fractures, cardiovascular disease and mortality. These observational studies have broadened the focus of CKD related mineral and bone disorders (MBDs) to include cardiovascular disease which is the leading cause of death in patients at all stages of CKD. All three of these processes (abnormal mineral metabolism, abnormal bone and extra skeletal calcification) are closely interrelated and together make a major contribution to the morbidity and mortality of patients with CKD. So we aim here to reveal the prevalence of markers of MBD in CKD stage 3-5 patients. Cardiovascular disease and mortality. These observational studies have broadened the focus of CKD related mineral and bone disorders (MBDs) to include cardiovascular disease which is the leading cause of death in patients at all stages of CKD. All three of these processes (abnormal mineral metabolism, abnormal bone and extra skeletal calcification) are closely interrelated and together make a major contribution to the morbidity and mortality of patients with CKD. So we aim here to reveal the prevalence of markers of MBD in CKD stage 3-5 patients. Results

### Subject and Methods

This prospective single centre study was conducted in Department of Medicine, Shri Ram Murti Smarak Institute of Medical Sciences, Bareilly - Nainital Rd, Bhoji Pura, Uttar Pradesh, India between October 2018 to December 2019. All CKD patients from stage 3- 5 were included. Likewise Patients having pre-existing systemic diseases like SLE/RA, liver disease, patients on steroids or other drugs which have effect on bone metabolism like calcium, phosphate binders, vit D, bisphosphonates, patients with primary bone diseases, patients on maintenance hemodialysis and h/o fracture in last 6 months are excluded from this study.

Glomerular filtration rate (eGFR) was estimated based on Cockcroft-Gault formula. Serum creatinine, albumin, calcium, phosphate (PO<sub>4</sub>), TAP, hemoglobin, uric acid, and urinary protein excretion were measured using standard laboratory techniques. Plasma intact parathormone (iPTH) was measured using the solid phase, two-site chemiluminescent enzyme-labeled immunometric assay.

Plasma 25-OH vitamin D (25-vitD) assay was done using the radioimmunometric assay. Radiological survey like lateral X-ray of skull, abdomen and 2D echocardiography was carried out to detect vascular and valvular calcification respectively. Statistical analyses are performed using SPSS software. Various parameters of the whole group were analyzed as well as the parameters were compared between the groups. Parametric variables were compared using unpaired t-test and Mann-Whitney Rank Sum test. Non-parametric variables were compared using Chi-square test. At P value<0.05 was taken as significant.

### Results

The total 175 cases, out of which 132(75.43 %) were males and 43(24.57 %) were females were taken in this hospital-based cross-sectional observational study. Male: female ratio is 3:1. Majority of patients were middle aged, i.e, in 41-60 years age group with mean age of 50.54 years, Out of all, 46 % patients were diabetic and 86 % patients were hypertensive.

**Table 1:** Distribution of calcium levels in different stage CKD patients.

	Calcium			Total
	<8.5	8.5-10.5	>10.5	
CKD 3 count	6	22	0	28
% within CKD 3	21.5%	78.5%	0%	100%
CKD 4 count	18	33	2	53
% within CKD4	33.9%	62.3%	3.8%	100%
CKD 5 count	46	48	0	94
% within CKD5	48.9%	51.1%	0%	100%
Total count	70	103	2	175
% within CKD	40.0%	58.8%	1.2%	100%

In stage 3 CKD patients (n=28), majority (78.5 %) had calcium levels in the normal range, only 21.5 % had calcium below normal range. In stage 4 patients (n=53), calcium levels were in the normal range in 62.3 % of patients, 33.9 % had calcium below normal range and 3.8 % had higher calcium levels than normal. In stage 5 CKD patients (94), calcium levels were in the normal range in 51.1 % of patients, while a significant number of patients (48.9 %) had low levels of calcium (Table 1). The results were found statistically significant with p value <0.05. Elevated phosphorus levels (>4.5mg/dl) in 13.8 % patients of CKD stage 3, 26.4 % patients of CKD stage 4 and 63.4 % patients of CKD stage 5 were found which is also statistically significant. [Table 2].

**Table 2:** Elevated phosphorus levels in 3 different stage CKD patients

	Phosphorus		Total
	2.5-4.5	4.5	
CKD 3 count	25	4	29
% within CKD3	86.2%	13.8%	100%
CKD 4	39	14	53
% within CKD 4	73.6%	26.4%	100%
CKD 5	34	59	93
% within CKD5	36.6%	63.4%	100%
Total count	98	77	175
% within CKD	56%	44%	100%

[Table 3] shows the results, which are found statistically significant with p value <0.05. The serum iPTH levels in CKD stage 3 patients, 51.9 % patients had normal iPTH levels whereas in significant number (48.1 %) of patients, iPTH was elevated above normal range. In CKD stage 4 patients, 66 % patients had iPTH above normal range and in only 34 % patients, iPTH was normal. In CKD stage 5 patients, iPTH was normal in only 7.4 % patients, 89.4% had high levels of iPTH and in 3.2% patients, it was below normal range.

**Table 3:** The serum iPTH level in CDK3, 4 and CDK5

	Intact parathyroid hormone		Total
	Normal	Above Normal	
CKD 3 count	0	14	27
% within CKD 3	0%	51.9%	100%
CKD 4 count	0	18	53
% within CKD 4	0%	34%	100%
CKD 5 count	3	7	95
% within CKD 5	3.2%	7.4%	100%
Total count	3	39	175
% within CKD	1.7%	22.3%	100%

Total alkaline phosphatase levels were within normal range in all patients of CKD stage 3. In stage 4 CKD patients, total ALP was within normal range in 94.3 % patients and remaining 5.7% had high levels. In stage 5 CKD patients, ALP was above normal in 45.3 % patients and normal in 54.7 % patients (Table 4). The results are statistically significant.

**Table 4:** Alkaline phosphatase levels in 3 different stages in CDK.

	<310	>310	Total
CKD 3 count	27	0	27
% within CKD 3	100%	0%	100%
CKD 4 count	50	3	53
% within CKD4	94.3%	5.7%	100%
CKD 5 count	52	43	95
% within CKD 5	54.7%	45.3%	100%
Total count	129	46	175
% within CKD	73.7%	26.3%	100%

Similarly (Table 5) shows 25 OH vitamin-D levels were below normal in 59.2% CKD stage 3 patients. Majority (70.3%) patients in CKD stage 4 had low levels of 25 OH vitamin-D and rest had normal levels. In CKD stage 5, 25 OH vitamin-D was low in 79.8 % patients and in rest, it was normal.


**Table 5:** 25 Hydroxy vitamin D in CDK3,4 and CDK5

	25 Hydroxy Vitamin D			Total
	<20	20 -30	>30	
CKD 3 count	8	8	11	27
% within CKD	29.6 %	29.6 %	40.8 %	100 %
CKD 4 count	20	18	16	54
% within CKD	37 %	33.9 %	29.7 %	100 %
CKD 5 count	45	30	19	94
% within CKD	47.9 %	31.9 %	20.2%	100 %
Total count	73	56	46	175
% within CKD	41.7 %	32 %	26.3 %	100 %

[Table 6] and [Figure 1] shows 8 patients of CKD stage 3, who have calcification. Among 21 patients of CKD stage 4, only 1(4.8 %) had evidence of vascular calcification and 10(29.4 %) had evidence of vascular calcification out of 34 patients of CKD stage 5 patients. The results were found statistically significant with p value <0.05.

**Table 6:** CKD Stage and Vascular Calcification.

	Vascular Calcification		Total
	Positive	Negative	
CKD 3	0 0 %	8 100 %	8 100 %
CKD 4	1 4.8 %	20 95.2 %	21 100 %
CKD 5	10 29.4 %	24 70.6 %	34 100 %
Total	11 19 %	52 81 %	63 100 %




**Fig 1:** CKD Stages and Vascular Calcification

Table 7 and Figure 2 shows 8 patients of CKD stage 3 who have undergone echocardiography, none had evidence of valvular calcification. Among 21 patients of CKD stage 4, all had evidence of valvular calcification and out of 34 patients of CKD stage 5 patients, 29(85.3 %) had evidence of valvular calcification.

**Table 7:** CKD stage and valvular calcification

	Valvular Calcification		Total
	Positive	Negative	
CKD 3	0 0 %	8 100 %	8 100 %
CKD 4	0 0 %	21 100 %	21 100 %
CKD 5	5 14.7 %	29 85.3 %	34 100 %
Total	5 7.9 %	58 92.1 %	63 100 %



**Fig 2:** CKD stages and valvular calcification

## Discussion

The mean age of our study population was higher (50.5 years) to other studies by Agarwal SK *et al.* (44 years), Sakuja V *et al.* (46.2 years) and B. Ghosh *et al.* (45.7 years) [5, 6, 7]. We observed that males outnumbered females (M: F = 3:1). There is male predominance among CKD population in most studies. In Nissenson's prevalence study from the United States, males had an overall prevalence of 1.6% and females 0.8%, this twofold ratio was maintained at all levels of serum creatinine [8]. Among Indian studies, Agarwal *et al.* [9]. Showed a male prevalence of 48% among patients with serum creatinine more than 1.8 mg/dl, while other hospital-based studies found males constituting 60–78% of CKD population [5, 6, 7]. One of the main reasons for these differences in may be that, in India, more males and younger persons attend hospitals than females and the elderly. MBDs are well described in patients with CKD.

Agarwal *et al.* described hypocalcemia in 29.9% and 49.6% in CKD stage 4 and 5, respectively, and hyperphosphatemia in 45% and 41.8%, respectively. LaClair *et al.* [10] found hypocalcemia in 8% and 28%, and hyperphosphatemia in 20% and 50% of patients of CKD stages 4 and 5, respectively. In our study, hypocalcemia was found in 21.5%, 33.9% and 48.9% cases of CKD stages 3, 4 and 5 respectively. Hyperphosphatemia was found in 13.8%, 26.4% and 63.4% cases of CKD stages 3, 4 and 5 respectively. Our study results corroborated with previous studies. Total ALP also signifies high turnover bone disease when elevated and interpreted in appropriate circumstances.

In this study, elevated ALP was present in 5.7% and 45.3% of patients of CKD stage 4 and 5, respectively. Indeed, KDIGO recommended that the treatment of MBD be based on trend in changes of biochemical parameters rather than on abnormalities at a single point of time. B. ghosh *et al.* found raised ALP in 43.59% and 76.66% of patients of CKD stage

4 and 5D, respectively. Jabbar, *et al.* [11] found raised bone alkaline phosphatase in 60% of their stage 4 and 5 CKD patients. Vitamin D abnormalities were common in all CKD stages. 60-80% patients had low levels of 25 hydroxy vitamin D. 1,25-dihydroxyvitamin D deficiency is known to occur during the progression of CKD, because the final hydroxylation step of 25-hydroxyvitamin D to 1, 25(OH) 2D is mediated by kidney 1 $\alpha$ -hydroxylase. Severity of deficiency did not correlate with CKD stage or other mineral abnormalities. In our study, the prevalence of deficiency of 25(OH)D3 increased as CKD progressed. Low

25(OH) D3 levels were found in 73.6 % of patients. Jabbar *et al.* [11] reported Vitamin D deficiency in 80%, and insufficiency in 13% of the patients. B. Ghosh *et al.* reported 83.13% of patients with CKD stage 4 and 5D had vitamin D level less than 30 ng/mL. Our study results corroborated with these studies. Literature shows that, hyperparathyroidism presents early in CKD & worsens with progression of CKD stages. There is an increase in the prevalence of hyperparathyroidism from CKD stage 4.

Hyperparathyroidism was present in 66% patients in CKD stage 4 & 89.4% patients in CKD stage 5 which was similar to Levin A *et al.* [12] study in which 56% patients in CKD with eGFR < 60ml/min had hyperparathyroidism. Agarwal *et al.* [5] found hyperparathyroidism in 57.8% of patients with CKD stage 4 and in 39.4% of patients with CKD stage 5. Jabbar, *et al.* [11].

Observed prevalence of hyperparathyroidism in 60% of their patients of CKD stage 4 and 5. In our study, hyperparathyroidism was higher than other studies.

Adynamic bone disease as evident by low iPTH levels was uncommon and found in 3.2 % patients of stage 5 CKD. Vascular (abdominal aortic) calcification was seen in 4.8 % and 29.4 % in CKD stage 4 and 5 patients respectively. Shantha *et al.* [13] using a lateral abdominal X-ray for screening, found a prevalence of 76.9% in 26 Indian pre-dialysis Stage 5 CKD patients who had a mean age of 56.6 years, 65% of whom were receiving calcium containing phosphate binders. A.T. Valson *et al.* [14] reported 6.8 % of cases having vascular calcification. Valvular calcification was present in 14.7 % of stage 5 CKD patients in our study, which is much lower than that reported in Caucasian pre-dialysis CKD subjects (31%) by Leskinen *et al.* [15]. Ghosh *et al.*, reported VC in 25% and 46% of Indian CKD Stage 4 and 5D patients respectively. A.T. Valson *et al.* [14] reported 96% of cases having valvular calcification. Among those having vascular or valvular calcification, most of the patients were older than 50 years and had evidence of hyperphosphatemia, hyperparathyroidism in comparison with those not having evidence of calcification.

## Conclusion

In conclusion, we observed Low 25 (OH) D levels, hyperparathyroidism, and hyperphosphatemia were the noticeable markers of CKD-MBD in our patients. Mineral bone disorder are common in CKD patients which start in early CKD stages & worsen with disease progression that causes morbidity and decreased quality of life. Hence, this

shows the importance of early recognition, understanding of their patho-physiological consequences & planning management strategies to prevent their progression, thereby reducing the cardiovascular morbidity & mortality.

**Source of Support:** Nil,

**Conflict of Interest:** None declared.

### References

1. Eknoyan G, Lameire N, Barsoum R, Eckardt KU, Levin A, Levin N *et al.* The burden of kidney disease: improving global outcomes. *Kidney international.* 2004; 66(4):1310-4.
2. Banerjee S, Ghosh B, Brojen T, Singh N, Singh S, Sharma OP *et al.* The high prevalence of chronic kidney disease-mineral bone disorders: a hospital-based cross-sectional study. *Indian J Nephrol.* 2012; 22:285-291.
3. National Kidney Foundation. K/DOQI clinical practice guidelines for chronic kidney disease: Evaluation, classification, and stratification. *Am J Kidney Dis.* 2002; 39(Suppl 1):S1-S266.
4. Martin KJ, Gonzalez EA. Strategies to minimize bone disease in renal failure. *Am J Kidney Dis.* 2001; 38:1430-1436.
5. Agarwal SK. Assessment of renal bone mineral disorder in naïve CKD patients: A single center prospective study. *Indian J Nephrol.* 2007; 17:96.
6. Sakuja V, Jha V, Ghosh AK, Ahmed S, Saha TK. Chronic renal failure in India. *Nephrol Dial Transplant.* 1994; 9:871-2.
7. Ghosh B, Brojen T. The high prevalence of chronic kidney disease mineral bone disorders: A hospital based cross sectional study. *Indian Journal of Nephrology.* 2012; 22:285-291.
8. Nissenson AR, Pereira BJ, Collins AJ, Steinberg EP. Prevalence and characteristics of individuals with chronic kidney disease in a large health maintenance organization. *Am J Kid Dis.* 2001; 37:1177-83.
9. Agarwal SK, Dash SC, Irshad M, Raju S, Singh R, Pandey RM. Prevalence of chronic renal failure in adults in Delhi, India. *Nephrol Dial Transplant.* 2005; 20:1638-42.
10. LaClair RE, Hellman RN, Karp SL, Kraus M, Ofner S, Li Q *et al.* Prevalence of calcidiol deficiency in CKD: A cross-sectional study across latitudes in the United States. *Am J Kidney Dis.* 2005; 45:1026-33.
11. Jabbar Z, Aggarwal PK, Chandel N, Khandelwal N, Sakuja V, Jha V *et al.* Noninvasive assessment of bone mineral status in indian CKD population- a cross-sectional study. *Indian J Nephrol.* 2007; 17:93.
12. Levin A, Bakris GL, Molitch M, Smulders M, Tian J, Williams LA *et al.* Prevalence of abnormal serum vitamin D, PTH, calcium, and phosphorus in patients with chronic kidney disease: results of the study to evaluate early kidney disease. *Kidney international.* 2007; 71(1):31-8.
13. Shantha GP, Kumar AA, Mancha A, Christopher M, Koshi R, Abraham G *et al.* Is abdominal aortic calcification score a cost-effective screening tool to predict atherosclerotic carotid plaque and cardiac valvular calcification in patients with end-stage renal disease *Indian J Nephrol.* 2012; 22:431-7.
14. Valson AT, Sundaram M, David VG, Deborah MN, Varughese S, Basu G *et al.* Profile of incident chronic kidney disease related-mineral bone disorders in chronic kidney disease Stage 4 and 5: A hospital based cross-sectional survey. *Indian journal of nephrology.* 2014; 24(2):97.
15. Leskinen Y, Paana T, Saha H, Groundstroem K, Lehtimäki T, Kilpinen S *et al.* Valvular calcification and its relationship to atherosclerosis in chronic kidney disease. *J Heart Valve Dis.* 2009; 18:429-38.