



E-ISSN: 2706-9575
P-ISSN: 2706-9567
IJARM 2021; 3(1): 275-277
Received: 15-12-2020
Accepted: 28-01-2021

Dr. Hemant Kumar Jain
Assistant Professor,
Department of General
Medicine, Government Medical
College Datia, Madhya
Pradesh, India

Dr. SN Shakya
PGMO, District Hospital
Datia, Madhya Pradesh, India

A study of prevalence of comorbidity among old age patients with hypertension

Dr. Hemant Kumar Jain and Dr. SN Shakya

DOI: <https://doi.org/10.22271/27069567.2021.v3.i1e.151>

Abstract

Background: The marked differences of prevalence in several studies are due to the different reference population, the various study designs, inclusion of diseases categories cardiovascular diseases in the older people is not seen in isolation, these groups may have other associated co-morbid conditions. The prevalence of comorbidity among people with hypertension is more common than those individuals with a normal BP. The management of multiple comorbidities requires more complex strategies to achieve effective care.

Objectives: To study of prevalence of comorbidity among old age patients with hypertension.

Material and Methods: All patients in our study above the age of 65 years, irrespective of their hypertensive status. All patients were classified according to blood pressure readings. All samples were venous blood gatherings in the morning after an 8 hr. fast. In all patients, ECG and fundus examination was done. 2D Echo was done whenever necessary.

Results: Of the 205 patients, all (100%) had a co-morbid illness present, of which ischemic heart disease contributed the maximum number of patients 58 (28.5%). This was followed by diabetes mellitus seen in 51 patients (25%). Other comorbidities included stroke in 15%, infections 14%, chronic obstructive pulmonary disease 13% and malignancy 4.5% patients.

Conclusion: Critical clinical examination and assessment of target organ damage, presence of co-morbid conditions in hypertensive individuals helps us in making the strategy for management.

Keywords: Old age, stroke, COPD, hypertension, ischemic heart disease, diabetes mellitus, co-morbidity

Introduction

The prevalence of comorbidity among people with hypertension is more common than those individuals with a normal BP. The comorbidity prevalence of hypertension has been estimated several times and estimates were inconsistent among publications from the government and those from academic societies. The marked differences of prevalence in several studies are due to the different reference population, the various study designs, inclusion of diseases categories, working definition of comorbidity, and representativeness of the data. Hypertension is a major health problem worldwide and its complications have significant socioeconomic impact.

Elderly persons with untreated hypertension are at higher risk of suffering from stroke and other major cardiovascular events ^[1]. As the population grows older, the incidence of hypertension, continues to increase in the developed and developing societies. Cardiovascular diseases in the older people is not seen in isolation, these groups may have other associated co-morbid conditions like arthritis, dementia, diabetes mellitus, dyslipidemia, vision disorder, ear nose and throat problems, orthopaedic problems, COPD and malignancy. Patients with multiple chronic conditions have on average a higher level of morbidity, poorer physical functioning and quality of life, a greater likelihood of persistent depression, and lower levels of social well-being ^[2-5]. Such patients incur increased risks of adverse drug events and mortality ^[6]. Despite the recent emphasis to conduct research on patients with multiple co-morbidities, even basic epidemiologic information such as prevalence is not well known. Thus, the present study was undertaken to study the prevalence of co-morbid conditions in elderly patients with hypertension. Among various routine data, information on several health statuses could be reflected better in health surveys than in medical records. Compared to other routine data, the survey, which includes both a social and health status, could provide better representative prevalence estimates of

Corresponding Author:
Dr. SN Shakya
PGMO, District Hospital
Datia, Madhya Pradesh, India

diseases, including comorbidity with hypertension.

Material and Methods

The present study was conducted in 205 patients above 65 years of age, who were admitted under various clinical departments like Medicine, Surgery, Gynaecology and allied branches in Government Medical College Datia, Madhya Pradesh. All patients were subjected to detailed clinical examination and investigations.

Inclusion criteria

All patients above the age of 65 years, irrespective of their hypertensive status, i.e., whether known hypertensive undergoing treatment, recently detected hypertensive or non-hypertensive.

Exclusion criteria

Patients below 65 years age group and with retroviral disease.

Methodology

The standard protocol was adapted when measuring for blood pressure (BP), blood sample collection and biochemical analysis. After of at least 5 minutes of rest and in both arms, supine as well as standing and in both lower limbs by a mercury sphygmomanometer. In patients who were bedridden, comatose, only supine blood pressure in arm and leg were taken. In patients with atrial fibrillation, a set of three readings and their mean was taken into consideration.

All patients were classified according to the VIIth US Joint National Committee on prevention, detection, evaluation and treatment of high blood pressure [7]. Hypertension was defined as an average SBP ≥ 140 mmHg, DBP ≥ 90 mmHg or the presence of antihypertensive agents. Prehypertension was defined as an average SBP 120-139 mmHg or DBP 80-89 mmHg. Diabetes mellitus (DM) was defined as having a fasting plasma glucose ≥ 126 mg/dL, the current use of anti-diabetic agents or the use of insulin prescribed by physician due to a previous diagnosis of diabetes. Impaired fasting glucose (IFG) was determined by the range; $100 \text{ mg/dL} \leq$ fasting plasma glucose (FPG) $< 126 \text{ mg/dL}$. Stroke, myocardial infarction (MI), angina, chronic kidney disease (CKD), and thyroid disease were defined as being a previous diagnosis given by a physician. CVD was defined as the previous diagnosis of stroke, MI, or angina. All samples were venous blood gatherings in the morning after an 8 hr. fast. In all patients, ECG and fundus examination was done. 2D Echo was done whenever necessary. Routine urine examination, blood urea, serum creatinine and USG abdomen was done to look for evidence of hypertensive nephropathy. CT Brain was done if there were clinical features of cerebrovascular accident to rule out haemorrhage and infarction.

Results

Out of these 205 patients, 131 (64%) were found to be hypertensive, of which 127 (62%) were male patients and 78 (38%) were female patients, as in Table 2. Of the 131 hypertensive patients, 76 (58%) were males and 55 (42%) were females. Out of the 131 patients, 82 (62.5%) were known hypertensive and 49 (37.5%) were newly detected. Only 17 patients had the blood pressure well controlled below 120/80 mm Hg. Of the 205 patients in our study,

dyslipidemia was found in 48 (23.5%) patients, of whom 29 (59.5%) were males and 19 (40.5%) were females. Isolated systolic hypertension is said to be present if the systolic blood pressure is more than 140 mm Hg and diastolic blood pressure is less than 90 mm Hg. Among 131 hypertensive patients, isolated systolic hypertension was diagnosed in 26 (19.9%) patients in whom 14 (52%) were males and 12 (48%) were females. Of the 205 patients taken up in our study, all (100%) had a co-morbid illness present, of which ischemic heart disease contributed the maximum number of patients (58 which accounted for 28.5%). This was followed by diabetes mellitus seen in 51 patients (25%). Other co-morbidities included stroke in 31 patients (15%), infections 29 (14%) patients, chronic obstructive pulmonary disease in 27 (13%) patients and malignancy was found in 9 (4.5%) patients. Table 1 and Figure 1.

Table 1: Co-morbid conditions in old age hypertensive patients

Co-Morbidity	No. of patients	Percentage
Ischemic heart disease	58	28.5%
Diabetes mellitus	51	25%
Stroke	31	15%
Infection	29	14%
COPD	27	13%
Malignancy	9	4.5%
Total	205	100%




Fig 1: Co-morbid conditions in hypertensive old age patients

Table 2: Sex ratio in our study

Sex	Number of cases	Percentage
Male	127	62%
Female	78	38%
Total	205	100%

Discussion

In agreement with similar studies, common comorbidities of hypertensive patients were dyslipidemia, obesity, and impaired fasting glucose. Seven of the eight chronic conditions were more prevalent in hypertensive patients than in adults without hypertension. The prevalence of comorbidity among people with hypertension is more common than those individuals with a normal BP. Co-morbidity is a common and notable status concerning the increasing complexity of care associated with it. It has been suggested that managing multiple comorbidities requires more complex strategies to achieve effective care. In our study, the incidence of hypertension was found to be 64% (131/205). In a study done by Farook *et al.*, [8] the incidence

of established hypertension among elderly was 61.4%. All the patients included in our study had a co-morbid illness present, of which ischemic heart disease contributed the maximum number of patients (58 which accounted for 28.5%). This was followed by diabetes mellitus seen in 51 patients (25%). Of the 58 patients detected for ischemic heart disease based on ECG, 75.5% (44 patients) had hypertension, while 24.5% (14 patients) were non hypertensive. In a study done by Dwivedi *et al.*, incidence of IHD detected was 57.6%^[9]. In Gupta *et al.* the incidence was as low as 3%^[10]. In the management of coronary artery disease among hypertensive, it was concluded by Bruce MP *et al.* that the use of short acting calcium channel blockers specially in high doses was associated with increased risk of myocardial infarction^[11] and the JNC VII has recommended diuretics and beta-blockers as the first line unless contraindicated^[3]. Low dose diuretics has been found to be safe and at the same time effective in the prevention of stroke, myocardial infarction, congestive cardiac failure and thus the total mortality. Diabetes mellitus is a widely accepted risk factor for IHD and stroke. In our study, DM was found in total of 51 patients (25%) of which 34 (17%) patients were hypertensive and 16 (8%) were non hypertensive. Farook *et al.* had the incidence of 37% in their study^[8]. Dwivedi *et al.* found the incidence to be 31.4%^[9] while Gupta *et al.* found the incidence of 13%^[10]. The findings in our study are in concordance with Dwivedi *et al.* The prevalence of DM is a significant predictor of a poor long term survival following strokes. Stroke is the second leading cause of death and disability in hypertensive patients^[8]. The systolic blood pressure is a better predictor of complications and borderline elevation of systolic blood pressure is associated with 42% increase in stroke and 52% increase in cardiovascular deaths. In our study, the incidence of stroke was 15% of which 13.5% were hypertensive and 5% were non hypertensive. Dwivedi *et al.* found 27.3% incidence in their study,^[9] Kulkarni *et al.* found the incidence of 15.4%^[12]. In an association analysis, the results of the present study are similar to the previous report. After adjustment for age, sex, marital status, socioeconomic status, insurance, and living area, hypertension had a positive association with the risk of multiple comorbidities. Stratified by sex among hypertensive patients, the risk for comorbidity was different without statistical significance. Only the risk for obesity was statistically significantly higher among males than among females.

Conclusion

Comorbidity is highly prevalent in Indian patients with hypertension. Critical clinical examination and assessment of target organ damage, presence of co-morbid conditions in hypertensive individuals helps us in making the strategy for management. The observations warn us to screen for hypertension in old age population at the early stage to prevent any complications.

References

1. Schwartz JB, Zipes DP. Cardiovascular disease in the elderly. In: Mann DL, Zipes DP, Libby PP, Bonow RO. Braunwald's Heart Disease: A Textbook of Cardiovascular Medicine. 10th edition. Philadelphia: Elsevier Science P1711-1741.
2. Bayliss EA, Ellis JL, Steiner JF. Barriers to self-

- management and quality-of-life outcomes in seniors with multimorbidities. *Ann Fam Med* 2007;5(5):395–402.
3. Fortin M, Lapointe L, Hudon C, Vanasse A, Ntutu AL, Maltais D. Multimorbidity and quality of life in primary care: a systematic review. *Health Qual Life Outcomes* 2004;2:51.
4. Fortin M, Bravo G, Hudon C, Lapointe L, Almirall J, Dubois MF. Relationship between multimorbidity and health-related quality of life of patients in primary care. *Qual Life Res* 2006;15(1):83–91.
5. Fortin M, Dubois MF, Hudon C, Soubhi H, Almirall J. Multimorbidity and quality of life: a closer look. *Health Qual Life Outcomes* 2007;5:52.
6. Gijzen R, Hoeymans N, Schellevis FG, Ruwaard D, Satariano WA, van den Bos GA. Causes and consequences of comorbidity: a review. *J Clin Epidemiol* 2001;54(7):661–674.
7. Chobanian AV, Bakris GL, Black HR, Cushman WC, Green LA, Izzo L Jr *et al.* National Heart, Lung, Blood Institute; National High Blood Pressure Education Program Coordinating Committee. Seventh report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure, Hypertension 2003;42;1206-1252.
8. Farook IF, Hussain S, Hasan M. Hypertension, Diabetes mellitus, hypercholesterolemia as risk factors of stroke. *Pakistan Journal of Med* 2003;42:412-418.
9. Dwivedi S, Singh G. Profile of hypertension in elderly subjects. *JAPI* 2000;48(2):1047-50.
10. Gupta HL, Yadav M *et al.* Study of prevalence of health problems in asymptomatic elderly individuals in Delhi. *JAPI* 2002;50:792-795.
11. Bruce MP, Heckbert SR, Koepsell JD *et al.* The risk of myocardial infarction associated with antihypertensive drug therapy. *JAMA* 1995;5:622-25.
12. Kulkarni V, Bhagat N, Hakim AV. Hypertension in the elderly. *JAPI* 2001;49:873-876.